

Bibliography

- Allison, P. D. (2000) Multiple imputation for missing data: a cautionary tale. *Sociological methods and Research*, 28, 301–309.
- Allison, P. D. (2002) *Missing Data*. Thousand Oaks, CA: Sage.
- Aylward, D. S., Anderson, R. A. and Nelson, T. D. (2010) Approaches to Handling Missing Data within Developmental and Behavioral Pediatric Research. *J. Developmental & Behavioral Pediatrics*. 31. 54-60.
- Barnard, J. and Meng, X.-L. (1999) Applications of multiple imputation in medical studies: from AIDS to NHANES. *Statistical Methods in Medical Research*, 8, 7–36.
- Burton, A. Altman, D. (2004) Missing Covariate Data within Cancer Prognostic Studies: a review of Current Reporting and Proposed Guidelines. *British J. Cancer*. 91, 4-8
- Carpenter, J., Pocock, S. and Lamm, C. J. (2002) Coping with missing data in clinical trials: a model based approach applied to asthma trials. *Statistics in Medicine*, 21, 1043–1066.
- Carpenter, J., Kenward, M., Evans, S. and White, I. (2004) Letter to the editor: Last observation carry forward and last observation analysis by J. Shao and B. Zhong, *Statistics in Medicine*, 22, 2429–2441. *Statistics in Medicine*, 23, 3241–3244.
- Carpenter, J. R., Kenward, M. G. and Vansteelandt, S. (2006) A comparison of multiple imputation and inverse probability weighting for analyses with missing data. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 169, 571-584.
- Carpenter, J. R., Kenward, M. G. (2013) *Multiple Imputation and its Applications*, Wiley N.Y.
- Casella, G. and George, E. I. (1992) Explaining the Gibbs sampler. *The American Statistician*, 46, 167–174.
- Chatterjee, N., Chen, Y. & Breslow N. E. (2003). A pseudoscore estimator for regression problems with two-phase sampling. *J. Amer. Statist. Ass.*, 98, 158–168.
- Chen, J. and Shao, J. (2000) Nearest neighbour imputation for survey data. *Journal of Official*

Statistics, 16, 113–131.

Cheung, Y. K. (2005) Exact two-sample inference with missing data *Biometrics*, 61, 524-531

Choi, S. C. and Lu, I. L. (1995) Effect of non-random missing data mechanisms in clinical trials. *Statistics in Medicine*, 14, 2675–2684.

Cook, R. J., Zeng, L. and Yi, G. Y. (2004) Marginal analysis of incomplete longitudinal binary data; a cautionary note on locf imputation. *Biometrics*, pp. 820–828.

Daniels, M.J. and Hogan, J.W. (2008) *Missing data in longitudinal studies; strategies for Bayesian modeling and sensitivity analysis*. Chapman and Hall (New York)

Dempster, A. P., Laird, N. M. and Rubin, D. B. (1977) Maximum likelihood from incomplete data via the em algorithm (with discussion). *Journal of the Royal Statistical Society Series B* (Statistical Methodology), 39, 1–38.

Diebolt, J. and Ip, E. H. S. (1996) Stochastic EM: method and application. In *Markov chain Monte-Carlo in practice* (Eds W. R. Gilks, S. Richardson and D. J. Spiegelhalter). London: Chapman and Hall.

Diggle, P. J. (1998) Dealing with missing values in longitudinal studies. In *Advances in the Statistical Analysis of Medical data* (Eds B. S. Everitt and G. Dunn), pp. 203–228. London: Arnold.

Diggle, P. J. and Kenward, M. G. (1994) Informative dropout in longitudinal data analysis (with discussion). *Journal of the Royal Statistical Society Series C (Applied Statistics)*, 43, 49–94.

Diggle, P.J., Patrick Heagerty, Kung-Yee Liang, Scott L Zeger (2002) *Analysis of Longitudinal Data* (Chapter 13 on Missing Values).

Egger, M., Davey Smith, G., Schneider, M., Minder, C. (1997) Bias in meta-analysis detected by a simple, graphical test, *British Medical Journal*, 315, 629-634.

Enders, C.K. (2010) *Applied Missing Data Analysis*, The Guilford Press (New York).

Engels, J. (2003) Imputation of missing longitudinal data: a comparison of methods. *Journal of Clinical Epidemiology*, 56, 968-976.

Fay, R. E. (1992) When are inferences from multiple imputation valid? *Proceedings of the Survey Research Methods Section of the American Statistical Association*, pp. 227–232.

- Fitzmaurice, G. and Laird, N. (1993) A likelihood-based method for analysing longitudinal binary response. *Biometrika*, 80, 141–151.
- Fitzmaurice, G., Laird, N. and Lipsitz, S. (1994) Analysing incomplete longitudinal binary responses: a likelihood-based approach. *Biometrics*, 50, 601–602.
- Fitzmaurice, G., Molenberghs, G. and Lipsitz, S. (1995) Regression models for longitudinal binary responses with informative dropout. *J. Royal Statistical Society Series B (Statistical Methodology)*, 57, 691–704.
- Follman, D. and Wu, M. (1995) An approximate generalized linear model with random effects for informative missing data. *Biometrics*, 51, 151–168.
- Forster, J. J. and Smith, P. W. F. (1998) Model-based inference for categorical survey data subject to non-ignorable non-response. *Journal of the Royal Statistical Society Series B (Statistical Methodology)*, 60, 57–60.
- Gad, A. M. and Abeer S. A. (2006) Analysis of longitudinal data with intermittent missing values using the stochastic EM algorithm, *Computational Statistics & Data Analysis*, 50, 2702-2714
- Gelman, A. and Raghunathan, T. E. (2001) Using conditional distributions for missing-data imputation, in discussion of 'using conditional distributions for missing-data imputation' by Arnold et al. *Statistical Science*, 3, 268–269.
- Gelman, A., Van Mechelen, I., Verbeke, G., Heitjan, D. F. and Meulders, M. (2005) Multiple imputation for model checking: Completed-data plots with missing and latent data *Biometrics*, 61, 74-85
- Gelman, A. and Rubin, D.B. (1992). Inference from iterative simulation using multiple sequences *Statistical Science*, 7, 457-511.
- Geyer, C.J. (1992) Practical Markov Chain Monte Carlo. *Statistical Science*, 7, 473-503
- Geweke, J. 1992. Evaluating the accuracy of sampling-based approaches to the calculation of posterior moments. pp. 169-93. In: *Bayesian Statistics 4* (eds J.M. Bernardo, J. Berger, A.P. Dawid and A.F.M. Smith.) Oxford University Press, Oxford.
- Gilks, W. R., Richardson, S. and Spiegelhalter, D. J. (1996) *Markov Chain Monte-Carlo in Practice*.

London: Chapman and Hall.

Glynn, R., Laird, N. and Rubin, D. (1986) Selection modelling versus mixture modelling with nonignorable nonresponse. In *Drawing Inferences from Self-selected Samples* (Ed. H. Wainer), pp. 115–142. New York: Springer-Verlag.

Godambe, V.P. (1960) An Optimum Property of Regular Maximum Likelihood Estimation. *Ann. Math. Statist.*, 31, 1208-1211.

Goetghebeur, E., Molenberghs, G. and Kenward, M. G. (1999) Sense and sensitivity when intended data are missing. *Kwantitatieve Technieken*, 62, 79–94.

González-Manteiga, W. and Pérez-González, A. (2004) Nonparametric mean estimation with missing data *Communications in Statistics: Theory and Methods*, 33, 277-303

Gould, A. (1980) A new approach to the analysis of clinical drug trials with withdrawals. *Biometrics*, 36, 721–727.

Greenland, S. (2005) Multiple-bias modelling for analysis of observational data (with discussion). *Journal of the Royal Statistical Society: Series A*, 168, 267–306.

Greenlees, W., Reece, J. and Ziexhang, K. (1982) Imputation of missing values when the probability of nonreponse depends on the variable being imputed. *J. Amer. Statist. Ass.*, 77, 251–256.

Han, P. and Wang, Lu (2013) Estimation with missing data: beyond double robustness. *Biometrika* 100, 417-430.

Hartley, H. and Hocking, R. (1971) The analysis of incomplete data. *Biometrics*, 27, 7783–808.

Heckman, J. J. (1976) The common structure of statistical models of truncation, sample selection and limited dependent variables and a simple estimator for such models. *Annals of Economic and Social Measurement*, 5, 475–492.

Hedeker, D. and Gibbons, R. D. (1997) Application of random-effects pattern-mixture models for missing data in longitudinal studies. *Psychological Methods*, 2, 64–78.

Heitjan, D. F. and Basu, A. (1996) Distinguishing ‘missing at random’ and ‘missing completely at random’. *J. Amer. Statist. Ass.*, 50, 207–213.

Heitjan D.F. and Little R.J.A. (1991). Multiple Imputation for the Fatal Accident Reporting System. *Journal of the Royal Statistical Society C*, 40(1), 13–29.

Heitjan, D. F. and Rubin, D. B. (1991) Ignorability and coarse data. *Annals of Statistics*, pp. 2244–2253.

Helms, R. W. (1992) Intentionally incomplete longitudinal designs: methodology and comparison of some full span designs. *Statistics in Medicine*, 11, 1889–1913.

Hershberger, Scott L. and Fisher, Dennis G. (2003) A note on determining the number of imputations for missing data. *Structural Equation Modeling*, 10, 648-650

Heyting, A., Tolboom, J. T. B. M. and Essers, J. G. A. (1992) Statistical Handling of Drop-Outs in Longitudinal Clinical Trials. *Statistics in Medicine*, 11, 2043–2061.

Hogan, J. and Laird, N. (1997a) Mixture models for joint distribution of repeated measures and event times. *Statistics in Medicine*, 16, 239–257.

Hogan, J. W. and Laird, N. M. (1997b) Mixture models for joint distribution of repeated measures and event times. *Statistics in Medicine*, 16, 239–257.

Hogan, J. W. and Laird, N. M. (1997c) Model-based Approaches to Analysing Incomplete Longitudinal and Failure Time Data. *Statistics in Medicine*, 16, 259–272.

Hollis, S. (2002) A graphical sensitivity analysis for clinical trials with non-ignorable missing binary outcome. *Statistics in Medicine*, 21, 3823–3834.

Horton, N. J. and Laird, N. M. (1999) Maximum likelihood analysis of generalized linear models with missing covariates. *Statistical Methods in Medical Research*, 8, 37–50.

Horton, Nicholas J. and Lipsitz, Stuart R. (2001) Multiple imputation in practice: Comparison of software packages for regression models with missing variables *The American Statistician*, 55, 244-254

Horton, N. J., Lipsitz, S. R. and Parzen, M. (2003) A potential for bias when rounding in multiple imputation. *The American Statistician*, 57, 229–232.

Horvitz, D. G. and Thompson, D. J. (1952) A generalisation of sampling without replacement from a

finite universe. *J. Amer. Statist. Ass.*, 47, 663–685.

Huang, L., Chen, M. and Ibrahim, J. G. (2005) Bayesian analysis for generalized linear models with nonignorably missing covariates. *Biometrics*, 61, 767–780.

Ibrahim, J. G., Chen, M. H. and Lipsitz, S. R. (1999) Monte-carlo em for missing covariates in parametric regression models. *Biometrics*, 55, 591– 596.

Ibrahim, J. G., Chen, M.-H. , Lipsitz, S. R. (2001) Missing responses in generalised linear mixed models when the missing data mechanism is nonignorable. *Biometrika*, 88, 551–564.

ICH E9 Expert Working Group (1999) Statistical Principles for Clinical Trials: ICH Harmonised Tripartite Guideline. *Statistics in Medicine*, 18, 1905–1942.

Imbens, G. W. and Rubin, D. B. (1997) Bayesian inference for causal effects in randomized experiments with noncompliance. *Annals of Statistics*, 25, 305–327.

Ip, E. H. S. (1994) A stochastic EM estimator in the presence of missing data. Technical report, Division of Biostatistics, Stanford University, Stanford, California, USA.

Ip, Edward H. (2002) On single versus multiple imputation for a class of stochastic algorithms estimating maximum likelihood. *Computational Statistics*, 17, 517-524

Jelicic, H., Phelps, E. and Lerner, R.M. (2009) Use of Missing Data Methods in Longitudinal Studies: The persistence of Bad Practices in Developmental Psychology. *Developmental Psychology*, 45, 1195-1199.

Kadane, J. B. (1993) Subjective Bayesian analysis for surveys with missing data. *Journal of the Royal Statistical Society, Series D (The Statistician)*, 42, 415–426.

Kalton, G. and Kasprzak, D. (1986) The treatment of missing survey data. *Survey Methodology*, 12, 1–16.

Kenward, M., Lesaffre, E. and Molenberghs, G. (1994) An application of maximum likelihood and generalized estimating equations to the analysis of ordinal data from a longitudinal study with cases missing at random. *Biometrics*, 50, 945–953.

Kenward, M. G. (1998) Selection Models for Repeated Measurements with Non-Random Dropout: An

Illustration of Sensitivity. *Statistics in Medicine*, 17, 2723–2732.

Kenward, M. G. and Molenberghs, G. (1998) Likelihood based frequentist inference when data are missing at random. *Statistical Science*, 13, 236– 247.

Kenward, M. G. and Molenberghs, G. (1999) Parametric models for incomplete continuous and categorical longitudinal data. *Statistical Methods in Medical Research*, 8, 51–83.

Kenward, M. G., Molenberghs, G. and Goetghebeur, E. (2001) Sensitivity analysis for incomplete categorical data. *Statistical Modelling*, 1, 31–48.

Kenward, M. G., Molenberghs, G. and Thijs, H. (2003) Pattern-mixture models with proper time dependence. *Biometrika*, 90, 53–71.

Kim, Jae Kwang (2004) Finite sample properties of multiple imputation estimators *The Annals of Statistics*, 32, 766-783.

King, G., Honaker, J., Joseph, A. and Scheve, K. (2001) Analysing incomplete political science data: an alternative algorithm for multiple imputation. *American Political Science Review*, 95, 49–69.

Klebanoff, M. A. and Cole, S. R. (2008). Use of Multiple Imputation in the Epidemiologic Literature. *American J. Epidemiology*, 168, 355-357.

Kofman, P. and Sharpe, I. G. (2003) Using multiple imputation in the analysis of incomplete observations in finance. *Journal of Financial Econometrics*, 1, 216-249.

Laird, N., Wu, M. et al. (1988) Analysis of data with missing values: discussion following meeting.

Laird, N. M. (1988) Missing data in longitudinal studies. *Statistics in Medicine*, 7, 305–315.

Landrum, M. B. and Becker, M. P. (2001) A multiple imputation strategy for incomplete longitudinal data, *Statistics in Medicine*, 20, 2741-2760.

Lasaffre, E., Molenberghs, G. and Dewulf, L. (1996) Effect of Dropouts in a Longitudinal Study: An Application of a Repeated Ordinal Model. *Statistics in Medicine*, 15, 1123–1141.

Lavori, P. W. (1992) Clinical trials in psychiatry: should protocol deviation censor patient data. *Neuropsychopharmacology*, 6, 39–48.

Lavori, P. W., Dawson, R. and Shera, D. (1995) A multiple imputation strategy for clinical trials with

- truncation of patient data. *Statistics in Medicine*, 14, 1913–1025.
- Lesaffre, E., Molenberghs, G. and Dewulf, L. (1996) Dropouts in a Longitudinal Study: An Application of a Repeated Ordinal Model. *Statistics in Medicine*, 15, 1123–1141.
- Li, K. H., Meng, X. L., Raghunathan, T. E. and Rubin, D. B. (1991a) Significance levels from repeated p-values with multiply imputed data. *Statistica Sinica*, 1, 65–92.
- Li, K. H., Raghunathan, T. E. and Rubin, D. B. (1991b) Large-sample significance levels from multiply-imputed data using moment-based statistics and an f references distribution. *J. Amer. Statist. Ass.*, 86, 1065–1073.
- Lindsey, J. K. (2000) Obtaining marginal estimates from conditional categorical repeated measurements models with missing data. *Statistics in Medicine*, 19, 801–809.
- Lipsitz, S., Parzen, M. and Ping Zhao, L. (2002) A degrees-of-freedom approximation in multiple imputation. *Journal of Statistical Computation and Simulation*, 72, 309–318.
- Little, R. (1986) A note about models for selectivity bias. *Econometrika*, 53, 1469–1474.
- Little, R. (1993a) Pattern-mixture models for multivariate incomplete data. *J. Amer. Statist. Ass.*, 88, 125–134.
- Little, R. and Wang, Y. (1996) Pattern-mixture models for multivariate incomplete data with covariates. *Biometrics*, 52, 98–111.
- Little, R. J. A. (1988) A Test of Missing Completely at Random for Multivariate Data With Missing Values. *J. Amer. Statist. Ass.*, 83, 1198–1202.
- Little, R. J. A. (1992) Regression with Missing X's: A Review. *J. Amer. Statist. Ass.*, 87, 1227–1237.
- Little, R. J. A. (1993b) Modelling the dropout mechanism in repeated-measures studies. *J. Amer. Statist. Ass.*, 88, 123–134.
- Little, R. J. A. (1994) A class of pattern-mixture models for multivariate incomplete data. *Biometrika*, 81, 471–483. 13
- Little, R. J. A. (1995) Modelling the drop-out mechanism in repeated measures studies. *J. Amer. Statist. Ass.*, 90, 1112– 1121.

- Little, R. J. A. and Rubin, D. B. (1987) *Statistical analysis with missing data*. Chichester: Wiley.
- Little, R. J. A. and Rubin, D. B. (1999) Contribution to the discussion of Scharfstein et al. *J. Amer. Statist. Ass.*, 94, 1096–1146.
- Little, R. J. A. and Rubin, D. B. (2002) *Statistical analysis with missing data* (second edition). Chichester: Wiley.
- Little, R. J. A. and Yau, L. (1996) Intent-to-treat analysis for longitudinal studies with drop-outs. *Biometrics*, 52, 471–483.
- Liu, C. (1999) Efficient ML estimation of the multivariate normal distribution from incomplete data. *J. Mult. Anal.* 69. 206-217.
- Louis, T. (1982) Finding the observed information matrix when using the EM algorithm. *J R Statist Soc B*, 44, 226–233.
- MacKinnon, A. (2010) The Use and Reporting of Multiple Imputation in Medical Research: A Review. *J. Internal Medicine* 268, 586-593
- Mallinckrodt, C., Sanger, T. M., Dubé, S., DeBrot, D. J., Molenberghs, G., Carroll, R. J., Potter, W. Z. and Tollefson, G. D. (2003a) Assessing and interpreting treatment effects in longitudinal clinical trials with missing data. *Biological Psychiatry*, 53, 754–760.
- Mallinckrodt, C., Scott-Clark, W. S., Carroll, R. J. and Molenberghs, G. (2003b) Assessing response profiles for incomplete longitudinal clinical trial data under regulatory considerations. *Journal of Biopharmaceutical Statistics*, 13, 179–190.
- Marschner, I. C. (2001) On stochastic versions of the EM algorithm. *Biometrika*, 88, 281–286.
- McCulloch, C. E. (1997) Maximum likelihood algorithms for generalized linear mixed models. *J. Amer. Statist. Ass.*, 92, 162–170.
- McLachlan, G. J. and Krishnan, T. (1997) *The EM algorithm and extensions*. Chichester: Wiley.
- McLeish, D.L. and Small, C.G. (1994) *Hilbert Space Methods in Probability and Statistical Inference*. J Wiley and Sons.
- McLeish, D.L. (2005) *Monte Carlo Methods and Finance*. J Wiley and Sons.

- McLeish, D.L. and Struthers, C. A. (2006) Estimation of regression parameters in missing data problems, *Can. J. Statist.* 34, 233-259.
- McLeish, D.L. (2002) Highs and Lows: Some Properties of the Extremes of a Diffusion and Applications in Finance. *Canadian Journal of Statist.* 30, 243-267
- Meng, X. L. (1994) Multiple-imputation inferences with uncongenial sources of input (with discussion). *Statistical Science*, 10, 538–573.
- Meng, X. L. and Rubin, D. B. (1992) Performing likelihood ratio tests with multiply-imputed data sets. *Biometrika*, 79, 103–111.
- Meng, X. L. and Romero, M. (2003) Discussion: Efficiency and self-efficiency with multiple imputation inference *International Statistical Review*, 71
- Michiels, B., Molenberghs, G. and Lipsitz, S. (1999) Selection models and pattern-mixture models for incomplete categorical data with covariates. *Biometrics*, 55, 978–983.
- Molenberghs, G. and Lesaffre, E. (1994) Marginal modelling of correlated ordinal data using a multivariate Plackett distribution. *J. Amer. Statist. Ass.*, 89, 633–644.
- Molenberghs, G., Kenward, M. and Lesaffre, E. (1997) The analysis of longitudinal ordinal data with non-random dropout. *Biometrika*, 84, 33–44.
- Molenberghs, G., Michiels, B., Kenward, M. G. and Diggle, P. J. (1998) Missing data mechanisms and pattern-mixture models. *Statistica Neerlandica*, 52, 153–161.
- Molenberghs, G., Goetghebeur, E., Lipsitz, S. R. and Kenward, M. G. (1999b) Non-random missingness in categorical data: strengths and limitations. *American Statistician*, 53, 110–118.
- Molenberghs, G., Goetghebeur, E., Lipsitz, S. R., Kenward, M. G., Lesaffre, E. and Michiels, B. (1999c) Missing data perspectives of the fluvoxamine data set: a review. *Statistics in Medicine*, 18, 2449–2464.
- Molenberghs, G., Kenward, M. G. and Goetghebeur, E. (2001a) Sensitivity analysis for incomplete contingency tables: the Slovenian plebiscite case. *Appl. Statist.*, 50, 15–29.
- Molenberghs, G., Verbeke, G., Thijs, H., Lesaffre, E. and Kenward, M. G. (2001b) Mastitis in dairy cattle: local influence to assess sensitivity of the dropout process. *Computational Statistics and Data*

Analysis, 37, 93–113.

Molenberghs, G., Thijs, H., Kenward, M. G. and Verbeke, G. (2002) How meaningful and sensitive are selection models and pattern-mixture models? *Proceedings of the XXIth International Biometric Conference, Invited Papers*, 295-315.

Molenberghs, G., Thijs, H., Jansen, I., Beunckens, C., Kenward, M. G., Mallinkrodt, C. and Carroll, R. J. (2004) Analyzing incomplete longitudinal clinical trial data. *Biostatistics*, 5, 445–464.

Molenberghs, G., Kenward, M. G. (2007) *Missing Data in Clinical Studies*, Wiley, New York.

More, M., Woolson, R. F. and Woodsworth, G. G. (1992) Application of empirical bayes inference to estimation of rate of change in the presence of informative right censoring. *Statistics in Medicine*, 11, 621–631.

Murray, G. and Findlay, J. G. (1988) Correcting for the Bias caused by Drop-outs in Hypertension Trials. *Statistics in Medicine*, 7, 941–946.

Nordheim, E. (1984) Inference from nonrandomly missing categorical data: an example from a genetic study of Turner's syndrome. *J. Amer. Statist. Ass.*, 79, 772–780.

Nan, B., Emond, M. J. and Wellner, J. A. (2004) Information bounds for Cox regression models with missing data. *The Annals of Statistics*, 32, 723-753.

Nielsen, Søren Feodor (2003) Proper and improper multiple imputation. *International Statistical Review*, 71, 593-607.

Orchard, T. and Woodbury, M. (1972) A missing information principle: theory and applications. In *Proceedings of the Sixth Berkely Symposium on Mathematics, Statistics and Probability*, Volume 1. (Eds L. M. L. Cam, J. Neyman and E. L. Scott), pp. 697–715. Berkeley: University of California Press.

Park, T. and Brown, M. B. (1994) Models for categorical data with nonignorable nonresponse. *J. Amer. Statist. Ass.*, 89, 44–52.

Peugh, J.L. and Enders, C.K. (2004) Missing Data in Educational Research: A Review of Reporting Practices and Suggestions for Improvement. *Review of Educational Research*, 74, 525-556.

- Pledge, G. and Hall, D. (1982) Correspondence on Gould, *Biometrics*, 1980, 36, 721–727. *Biometrics*, 28, 276–278.
- Pledger, G., Hall, D. and Gould, A. L. (1982) Withdrawals from Drug Trials. *Biometrics*, pp. 276–278.
- Raab, G. M. and Donnelly, C. A. (1999) Information on sexual behaviour when some data are missing. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 48, 117–133.
- Raboud, J. M., Montaner, J. S. G., Thorne, A., Singer, J., Schechter, M. T. and Canadian HIV trials group (1996) Impact of Missing Data Due to Dropouts on Estimates of the Treatment Effect in a Randomized Trial of Antiretroviral Therapy for HIV-Infected Individuals. *Journal of AIDS and Human Retrovirology*, 12, 46–55.
- Raghunathan, T. E., Lepkowski, J. M., Van Hoewyk, J. and Solenberger, P. (2001) A multivariate technique for multiply imputing missing values using a sequence of regression models. *Survey Methodology*, 27, 85–95.
- Rao, C.R. (1973) *Linear Statistical Inference and its Applications*. Wiley.
- Rao, J. N. K. (1996) On variance estimation with imputed survey data. *J. Amer. Statist. Ass.*, 91, 499–506.
- Robbins, H., Monro, S. (1951) A Stochastic Approximation Method. *Ann. Math. Statist.*, 22, 400–407.
- Robins, J. and Rotnitzky, A. (1995a) Semi-parametric estimation of models for the means and covariances in the presence of missing data. *Scandinavian Journal of Statistics*, 22, 323–333.
- Robins, J. M. and Rotnitzky, A. (1995b) Semiparametric efficiency in multivariate regression models with missing data. *J. Amer. Statist. Ass.*, 90, 122–129.
- Robins, J. M. and Wang, N. (2000) Inference for imputation estimators. *Biometrika*, 85, 113–124.
- Robins, J. M., Rotnitzky, A. and Zhao, L. P. (1994) Estimation of regression coefficients when some regressors are not always observed. *J. Amer. Statist. Ass.*, 89, 846–866.
- Robins, J. M., Rotnitzky, A. and Zhao, L. P. (1995a) Estimation of regression coefficients when some regressors are not always observed. *J. Amer. Statist. Ass.*, 89, 846–866.
- Robins, J. M., Rotnitzky, A. and Zhao, L. P. (1995b) Analysis of semiparametric regression models for

- repeated outcomes in the presence of missing data. *J. Amer Statist. Ass.*, 90, 106–129.
- Robins, J. M., Rotnitzky, A. and Scharfstein, D. O. (1998) Semiparametric regression for repeated outcomes with non-ignorable non-response. *J. Amer. Statist. Ass.*, 93, 1321–1339.
- Rotnitzky, A. and Robins, J. M. (1997) Analysis of semi-parametric regression models with nonignorable nonresponse. *Statistics in Medicine*, 16, 81–102.
- Rotnitzky, A., Robins, J. M. and Scharfstein, D. O. (1998) Semiparametric regression for repeated outcomes with nonignorable nonresponse. *J. Amer. Statist. Ass.*, 93, 1321–1339.
- Royston, P. (2004) Multiple imputation of missing values. *The Stata Journal*, 3, 227–241.
- Rubin, D. (1996) Multiple imputation after 18 years. *J. Amer. Statist. Ass.*, 91, 473–490.
- Rubin, D. B. (1976) Inference and missing data. *Biometrika*, 63, 581–592.
- Rubin, D. B. (1987) *Multiple imputation for nonresponse in surveys*. New York: Wiley.
- Saltelli, A., Chan, K. and Scott, E. M. (eds) (2000) *Sensitivity analysis*. Chichester: Wiley.
- Sarndal, C. E., Swensson, B. and Wretman, J. (1992) *Model assisted Survey Sampling*. New York: Springer.
- Schafer, J. L. (1997) *Analysis of incomplete multivariate data*. London: Chapman and Hall.
- Schafer, J. L. (1999) Multiple imputation: a primer. *Statistical Methods in Medical Research*, 8, 3–15.
- Schafer, J. L. (2001) Multiple imputation with pan. In *New Methods for the Analysis of Change* (Eds L. M. Collins and A. G. Sayer), pp. 355–377. Washington, DC: American Psychological Association.
- Schafer, Joseph and Harel, Ofer (2002) Two-stage multiple imputation *ASA Proceedings of the Joint Statistical Meetings*, 1359-1363.
- Schafer, J. L. (2003) Multiple imputation in Multivariate Problems when the imputation and analysis models differ. *Stat. Neerlandica*, 57, 19-35.
- Schafer, J. L. and Graham, J. W. (2002) Missing data: our view of the state of the art. *Psychological Methods*, 7, 147–177.

- Schafer, J. L. and Yucel, R. (2002) Computational strategies for multivariate linear mixed-effects models with missing values. *Journal of Computational and Graphical Statistics*, 11, 421–442.
- Scharfstein, D. O. and Irizarry, R. A. (2003) Generalized additive selection models for the analysis of studies with potentially nonignorable missing outcome data. *Biometrics*, 59, 601–613.
- Scharfstein, D. O., Rotnitzky, A. and Robins, J. M. (1999) Adjusting for nonignorable drop-out using semi-parametric nonresponse models (with comments). *J. Amer. Statist. Ass.*, 94, 1096–1146.
- Schenker N. and Taylor J.M.G. (1996). Partially Parametric Techniques for Multiple Imputation. *Computational Statistics & Data Analysis*, 22, 425–446.
- Schluchter, M. D. (1992) Methods for the analysis of informatively censored longitudinal data. *Statistics in Medicine*, 11, 1861–1870.
- Shao, J. and Zhong, B. (2003) Last observation carry-forward and last observation analysis. *Statistics in Medicine*, 22, 3241–3244.
- Scheuren, Fritz (2005) Multiple imputation: How it began and continues. *The American Statistician*, 59, 315-319.
- Shi, J. Q. and Copas, J. B. (2002) Publication bias and meta-analysis for 2x2 tables: an average markov chain monte carlo em algorithm. *Journal of the Royal Statistical Society Series B (Statistical Methodology)*, 64, 221–236.
- Shih, W. J. and Quan, H. (1997) Testing for Treatment Differences with Dropouts Present in Clinical Trials -a Composite Approach. *Statistics in Medicine*, 16, 1225–1239.
- Sinharay, S., Stern, H. S. and Russell, D. (2001) The use of multiple imputation for the analysis of missing data. *Psychological Methods*, 6, 317–329.
- Skinner, C. and Rao, J. N. K. (2002) Jack-knife variance estimation for multivariate statistics under hot deck imputation from common donors. *Journal of Statistical Planning and Inference*, 102, 421–422.
- Stasny (1986) Estimating gross flows using panel data with nonresponse: an example from the Canadian Labour Forces Survey. *J. Amer. Statist. Ass.*, 81, 42–47.
- Struthers, C. and McLeish, D.L. (2011) A Particular Diffusion Model for Incomplete Longitudinal Data:

Application to CD4 Counts and Time-to-Event Data in the Multicenter AIDS Cohort Study. *Biostatistics* 12 (3): 493-505.

Su, Y.S., Gelman, A., Hill, J.L. and Yajimi, M. (2011) Multiple imputation with diagnostics (mi) in R: Opening windows into the black box. *J. Stat. Software.*, 45.

Sui, D. and Fouladi, R. T. (2003) Missing data in a small longitudinal study: A comparison of monotone, propensity, or non-monotone multiple imputation methods with complete case and general linear mixed model analyses. *ASA Proceedings of the Joint Statistical Meetings*, 4134-4140 *American Statistical Association (Alexandria, VA)*

Tang, G., Little, R. J. A. and Raghunathan, T. E. (2003) Analysis of multivariate missing data with nonignorable nonresponse. *Biometrika*, 90, 747-764.

Tanner, M. and Wong, W. (1987) The calculation of posterior distributions by Data Augmentation (with discussion). *J. Amer. Statist. Ass.*, 82, 528–550.

Taylor, J. M. G., Cooper, K. L., Wei, J. T., Sarma, R. V., Raghunathan, T. E. and Heeringa, S. G. (2002) Use of Multiple Imputation to Correct for Nonresponse Bias in a Survey of Urologic Symptoms among African-American Men. *American Journal of Epidemiology*, 156, 774–782.

Thijs, H., Molenberghs, G. and Verbeke, G. (2000) The milk protein trial: influence analysis of the dropout process. *Biometrical Journal*, 5, 617–646.

Thijs, H., Molenberghs, G., Michiels, B., Verbeke, G. and Curran, D. (2002) Strategies to fit pattern-mixture models. *Biostatistics*, 3, 245–265.

Touloumi, G., Pocock, S. J., Babiker, A. G. and Darbyshire, J. H. (1999) Estimation and comparison of rates of change in longitudinal studies with informative dropout. *Statistics in Medicine*, 18, 1215–1233.

Touloumi, G., Babiker, A. G., Kenward, M. G., Pocock, S. J. and Darbyshire, J. H. (2003) A comparison of two methods for the estimation of precision with incomplete longitudinal data, jointly modelled with a time-to-event outcome. *Statistics in Medicine*, 22, 3161–3175.

Toutenburg, H. and Srivastava, V. K. (2002) A revisit to the application of weighted mixed regression estimation in linear models with missing data. *American Journal of Mathematical and Management Sciences*, 22, 281-301.

- Troxel, A., Harrington, D. and Lipsitz, S. (1998a) Analysis of longitudinal data with non-ignorable non-monotone missing values. *Journal of the Royal Statistical Society, Series C, (Applied Statistics)*, 47, 425–438.
- Troxel, A., Lipsitz, S. and Harrington, D. (1998b) Marginal models for the analysis of longitudinal data with nonignorable non-monotone missing data. *Biometrika*, 85, 661–673.
- Vach, W. and Blettner, M. (1991) Biased Estimation of the Odds Ratio in Case-Control Studies due to the Use of Ad Hoc methods of Correcting for Missing Values for Confounding Variables. *American Journal of Epidemiology*, 134, 895–907.
- Vach, W. and Blettner, M. (1998) Missing Data in Epidemiologic Studies. In *Encyclopedia of Biostatistics* (Eds P. Armitage and T. Colton), volume 4, pp. 2641–2654. John Wiley and Sons, Chichester, United Kingdom.
- Vach, W. and Blettner, M. (1999) Logistic regression with incompletely observed categorical covariates — investigating sensitivity against violation of the Missing at Random assumption. *Statistics in Medicine*, 54, 1315– 1329.
- Van Buuren, S., Boshuizen, H. C. and Knook, D. L. (1999) Multiple imputation of missing blood pressure covariates in survival analysis. *Statistics in Medicine*, pp. 681–694.
- Van Buuren, S. (2012) *Flexible imputation of missing data*, CRC Press (Bocaraton)
- van Buuren, S. and K. Groothuis-Oudshoorn, K. (2011) MICE: Multivariate imputation by chained equations in R. *J. Stat. Software.*, 45, 1–67
- Van der Laan, M. J. and Robins, J. (2003) Unified methods for censored longitudinal data and causality. New York: Springer.
- Vansteelandt, S. and Goetghebeur, E. (2001) Analysing the sensitivity of generalized linear models to incomplete outcomes via the ide algorithm. *Journal of Graphical and Computational Statistics*, 10, 656–672.
- Verbeke, G. and Lesaffre, E. (1999) The effect of drop-out on the efficiency of longitudinal experiments. *Journal of the Royal Statistical Society Series C (Applied Statistics)*, 48, 363–375.
- Verbeke, G. and Molenberghs, G. (2000) *Linear Mixed Models for Longitudinal Data*. New York:

Springer Verlag.

- Verbeke, G., Molenberghs, G., Thijs, H., Lesaffre, E. and Kenward, M. G. (2001) Sensitivity analysis for non-random dropout: a local influence approach. *Biometrics*, 57, 7–14.
- Verzilli, C. and Carpenter, J. R. (2002a) A Monte Carlo EM algorithm for random-coefficient-based dropout models. *Journal of Applied Statistics*, 29, 1011–1021.
- Verzilli, C. and Carpenter, J. R. (2002b) Assessing uncertainty about parameter estimates with incomplete repeated ordinal data. *Statistical Modelling*, 2, 203–215.
- Von Hippel, P.T. (2015) New Confidence intervals and bias comparisons show that maximum likelihood can beat multiple imputation in small samples. (to appear: Structural Equations Modeling)
- Wainer, H. (ed.) (1986) *Drawing Inferences from Self-selected Samples*. New-York: Springer-Verlag.
- Wang, N. and Robins, J. M. (1998) Large-sample theory for parametric multiple imputation procedures. *Biometrika*, 85, 935–948.
- Wang-Clow, F., Lange, M., Laird, N. M. and Ware, J. H. (1995) A simulation study of estimators for rates of change in longitudinal studies with attrition. *Statistics in Medicine*, 14, 283–297.
- Ware, J.H. , Harrington, D., Hunter, D.J. , Agostino, R. B. (2012) Missing Data. *New England J. Med.* 1353-1390
- Wei, G. C. G. and Tanner, M. A. (1990) A Monte Carlo Implementation of the EM Algorithm and the Poor Man's Data Augmentation Algorithms. *J. Amer. Statist. Ass.*, 85, 699–704.
- Wei, L. J. and Johnson, W. E. (1985) Combining dependent tests with incomplete repeated measurements. *Biometrika*, 72, 359–364.
- White, I., Carpenter, J., Evans, S. and Schroter, S. (2004) Eliciting and using expert opinions about non-response bias in randomised controlled trials. Technical report, Medical Statistics Unit, London School of Hygiene and Tropical Medicine.
- White, I. R., Moodie, E., Thompson, S. G. and Croudace, T. (2003) A modelling strategy for the analysis of clinical trials with partly missing longitudinal data. *int J Meth Psych Res*, pp. 139–150.
- Whitehead J. (1997) *The design and analysis of sequential clinical trials*. Chichester, Wiley (1997)

- Wood, A. M., White, I. R. and Thompson, S. G. (2004) Are missing outcome data adequately handled? A review of published randomized controlled trials in major medical journals. *Clinical Trials*, 1, 368–376.
- Wu, M. and Bailey, K. (1989) estimation and comparison of changes in the presence of informative right censoring: conditional linear model. *BCS*, 45, 939–955.
- Wu, M. C. and Bailey, K. (1988) Analysing changes in the presence of informative right censoring caused by death and withdrawal. *Statistics in Medicine*, 7, 337–346.
- Wu, M. C. and Bailey, K. R. (1998) Analysing changes in the presence of informative right censoring by modelling the censoring process. *Statistics in Medicine*, 7, 175–188.
- Wu, M. C. and Carroll, R. J. (1988) Estimation and comparison of changes in the presence of informative right censoring by modelling the censoring process. *Biometrics*, 44, 175–188.
- Yau, L. H. and Little, R. J. A. (2001) Inference for the complier-average causal effect from longitudinal data subject to noncompliance and missing data, with application to a job training assessment for the unemployed. *J. Amer. Statist. Ass.*, 96, 1232–1244.
- Zhang, Paul (2003) Multiple imputation: Theory and method. *International Statistical Review*, 71, 581–592.
- Zhao, Y., Lawless, J. and McLeish, D.L. (2009) Likelihood Methods for Regression Models with Expensive Variables Missing by Design. *Biometrical Journal*, 51, 1–14.